



International Journal of Engineering Research and Science & Technology

ISSN : 2319-5991
Vol. 6, No. 1
February 2017



www.ijerst.com

Email: editorijerst@gmail.com or editor@ijerst.com

Research Paper

EFFECTIVENESS OF BIOLOGICAL ORGANIC FERTILIZER PLUS INORGANIC FERTILIZER ON THE GROWTH AND PRODUCTION PLANT VARIETY SOME HYBRID CORN (*ZEA MAYS* L.) AND WEEDS IN LOWLAND SWAMP AREA

Rosmiah¹, Gusmiatun¹, I in Siti Aminah¹, Minwal¹ and Neni Marlina^{2*}*Corresponding Author: Neni Marlina ✉ marlina002@yahoo.com

Increased food production is currently directed at suboptimal as lowland swamp that is widespread in several regions in Indonesia (Purwanto, 2005; and Subagyo, 2006). The main problem in the utilization of lowland swamp that waterlogging, soil acidity, the presence of Al and Fe cations binding phosphorus and nutrient-poor (Alihamsyah and Ar-Riza, 2006). Efforts to improve the productivity of maize in lowland swamp area through organic fertilizer enriched with microorganisms called bio-organic fertilizer (bio-fertilizer) and the use of inorganic fertilizers with low dose. The research aims to identify and determine the dose of organic fertilizer plus best inorganic fertilizer on the growth and production of several varieties of hybrid corn plants (*Zea mays* L.) and weeds in lowland swamp area. Field trial was conducted from April 2016 to August 2016, in the village of Pulau Semambu Ogan Ilir, South Sumatra Province. Statistical experiments using a randomized block design factorial with 15 combinations of treatments and 3 replications. Factor I: Biological Organic Fertilizer dose (BOF) + Inorganic Fertilizers (T) consisting of T0 = BOF + 0% N, P, K; T1 = BOF + 25% N, P, K; T2 = BOF + 50% N, P, K; T3 = BOF + 75% N, P, K; T4 = BOF + 100% N, P, K. Factor II: hybrid corn varieties (V), namely Pioneer V1 = 27, V2 = Bisi 18, V3 = DK 85. Variables observed that High Plant (cm) Number of leaves (leaf), length of cob (cm), diameter of cob (cm), weight of 100 seeds (g), shelled weight per plot (kg), plant remains dry weight (g). The number and type Weeds (SDR), Weeds wet weight (g), Weeds Dry Weight (g). Result: The biological treatment of organic fertilizer + 75% N, P, K is the best treatment shelled weight per plot of 714.67 kg. There were changes in weed populations dominated by *Imperata cylindrica* before treatment and after the study *Cyperus compressus* weeds dominated.

Keywords: Biological organic fertilizer dose, Inorganic fertilizer, Hybrid corn, Weeds, Lowland swamp area

¹ Lecturer of Agrotechnology Departemen Agriculture Faculty of University of Muhammadiyah Palembang A. Yani Street, 13 Ulu Palembang.

² Lecture of Agrotechnology Departemen Agriculture Faculty of University of Palembang.

INTRODUCTION

Maize (*Zea mays* L.) is the second staple crop after rice and used as food sources of carbohydrates. The development of farming systems and agribusiness corn in swamp land shown good prospects for still vast swamp land in Indonesia that has not been earned.

The main problem in the utilization of swampy wetlands that waterlogging, soil acidity, the presence of Al and Fe cations binding phosphorus and nutrient-poor (Alihamsyah and Ar-Riza, 2006). Therefore, support cultivation technology and management of land, water, and nutrients in corn development in wetlands indispensable includes land preparation and tillage, seeding, use of improved varieties, amelioration and fertilization, mulching and organic ingredients.

Organic fertilizers are essential components in organic farming that play a role in maintaining soil fertility and quality of crops through improved biological activities that ultimately can interact with the physical properties and chemical soil (Mahdil and Hassan, 2010).

Applications include livestock manure organic fertilizer, biochar, husk, rice straw or a combination thereof can be used to enhance the ability of the corn crop in the face of poor soil conditions Iqbal (2008), stating that the application of biological fertilizer on maize is able to increase an average of 50-97% uptake of macro and micro nutrients from 58.9 to 63.2%.

Organic fertilizers such as manure chicken manure can decompose quickly if aided by microbes include *Lactobacillus* lactic acid bacteria, photosynthetic bacteria and *Streptomyces* sp. and yeasts. These microbes provide a good influence on the quality of organic fertilizer, while the availability of nutrients in the

fertilizer was influenced by the length of incubation time needed by bacteria to degrade organic fertilizer. The results showed that the fertilizer that has been composted and incubated for 20 days provide a real influence on the growth of maize (Siburán, 2007). Besides the use of organic fertilizers, the addition of inorganic fertilizer needs to be done to improve the growth of corn plants but pursued with lower doses than the recommended, it is because in organoc fertilizer, have nutrient elements more fully and quickly available to plants, while organic fertilizers are Slow release.

Additionally weeds become one of the obstacles in swampy areas. Weeds are plants that harm other plants that are nearby or staple crops and grow in unwanted places. Due to the nature of the harm, then wherever weeds grow always be revoked, daytime, and even burned. (Suryaningsih, 2011).

The purpose of this study is to find out and get a dose of organic fertilizer plus N, P, K and best varieties on the growth and yield of hybrid maize (*Zea mays* L.) as well as the growth of the weed population in lowland swamp area.

MATERIALS AND METHODS

Research conducted at the Experimental Garden, Faculty of Agriculture in Campus C Muhammadiyah University Palembang Pulau Semambu Village, District of Indralaya Utara, Ogan Ilir Regency, in South Sumatra, from April to August 2016. Materials used in the research are agricultural lime, hybrid corn seeds (Pioneer 27, Bisi 18 and DK 85), chicken manure organic fertilizer, biological fertilizer, rice husk, bran, sugar, Urea, SP 36, KCl, Furadan and Decis.

This study uses a randomized block design with 15 combinations of treatments and 3

replications. A spacing of 70 x 25 cm corn planted in plots with 45 plots. Planting corn seeds 2 points per planting hole with ditugal, cultivation techniques and maintenance refers to the integrated crop management unless weeds were allowed to grow. Initial soil analysis is done based on the results of laboratory analysis P.T Binasawit Makmur, *Soil and Plant Tissue Analysis Laboratory Laboratoryegrated*, Palembang, South Sumatra.

To determine the effect among the treatments performed by analysis of variance followed by SAS program HSD test level of 5%. Variables measured include: plant height, number of leaves, cob length, cob diameter, weight of 100 seeds, shelled weight and dry weight per plot of crop residu. Observations weeds are the type and amount of weeds (SDR), fresh weight and dry weight of weeds.

RESULTS

The soil used in this study have the chemical and physical properties as listed in Table 1.

Based on the assessment criteria for the chemical and physical soil properties according Hardjowigeno (1995), the soil used in this study included the category of low-fertility soils with pH H₂O as very acid. Subagyo (2006), that the swampy soil pH ranging from 4.0 to 5.5 and the content of macro nutrients are low. The results of analysis of soil on top, despite the availability of nutrients P high, but many sequestered metal ions in the soil such as aluminum, to form Al-P and may cause P can not be absorbed by plants. Attempts to improve soil fertility in the study were given agricultural lime 5 tons/ha, aims to increase the pH and the addition of organic fertilizer and inorganic fertilizer for plants that grow on it can grow and produce the maximum. Organic

Table 1: Characteristics of the Soil Before Planting in Soil Research in Pulau Semambu Village, 2016

No.	Observations	Results	Criteria
1	pH H ₂ O	4,17	Very Acid
2	pH KCl	3,99	
3	N-total (%)	0,48	Average
4	C-organic (%)	14,32	Very High
5	KTK (me/100 g)	24,76	Average
6	K-dd (me/100 g)	0,23	Low
7	Ca-dd (me/100 g)	0,07	Very Low
8	Mg-dd (me/100 g)	0,05	Very Low
9	Na-dd (me/100 g)	0,05	Very Low
10	P ₂ O ₅ (me/100 g)	127,59	Very High
11	K ₂ O (me/100 g)	7,19	Very Low
12	P Bray I (ppm)	8,37	Average
13	Al ₂ O ₃ (%)	6,58	
14	Soil Texture		Clay Sandy
	Sand (%)	70,00	
	Dust (%)	14,00	
	Clay (%)	16,00	

Note: Description: Criteria Land Research Center (1983) and the Soil Research Institute (2005).

fertilizers are used in this study has been added to the biological fertilizer and incubated for 20 days (on a preliminary study with a C/N ratio is 19.19). According Hardjowigeno (1995), that the best organic fertilizers have a value of C/N-ratio range from 10-20.

Results of analysis of variance Table 2 shows that treatment of organic biological fertilizer plus N, P, K and interaction effect was not significant to all of the observed variables Treatment varieties significant to very significant to all of the observed variables, but no real effect on the cob length and weight of 100 seeds.

Table 2: Results of Analysis of Variance Influence Biological Treatment of Organic Fertilizer Plus N, P, K and Varieties of the Observed Variables

Treatment Parameters	Observed			Diversity Coefficient (%)
	T	V	I	
Plant Height (cm)	ns	*	ns	32,01
Number of leaves (sheet)	ns	*	ns	17,05
Cob Length (cm)	ns	ns	ns	20,48
Cob Diameter tongkol (cm)	ns	*	ns	15,47
Weight of 100 seeds (g)	ns	ns	ns	10,87
Seed Weight per plot (g)	ns	**	ns	25,36
Dry Weight of crop Residue (g)	ns	**	ns	27,21

Note: Description: ns = non significant; * = significant; ** = very significant; T = organic biological fertilizer plus organic fertilizer; V = varieties; I = interaction.

Data influence biological treatment of organic fertilizer plus N, P, K of the variables of growth and production of hybrid corn and its interactions shown in Table 3 and Table 5. The effect of treatment of different varieties manifest until very real, then continued test of HSD in Table 4.

Table 3 shows that the biological treatment of organic fertilizer plus N, P, K can promote the

growth and production of corn increased with increasing doses of fertilizers applied and the best at the lowest T0 and T3 treatment.

Table 4 shows that treatment of different varieties manifest until very significant to the growth of corn and variables positively correlated with yield components, it is best to V₁ (varieties Pioneer 27), V₂ (Bisi 18) and V₃ (DK 85).

To see the relationship between plant height with a weight of seed per plot shown in Figure 1, and the relationship between plant height with a dry weight of crop residue shown in Figure 2.

Figure 1 shows that the higher the dose of fertilizer given a high increase in corn and positively correlated with the weight of shelled corn kernels per plot to the extent of fertilizer combination BOF + 75% N, P, K. Similarly, what happened in Figure 2 shows that the increase in plant height due to the increase in dry weight of crop residue of corn and peak in the combination treatment of manure BOF + 75% N, P, K.

Nutrient Uptake of N, P dan K in Plant Tissue (%)

Nutrient Uptake of N, P and K Plant Tissue (%) Uptake of N, P and K corn plant tissue performed at maximum vegetative phase (8

Table 3: Effect of Biological Organic Fertilizer (BOF) Plus N, P, K of the Variables of Growth and Production of Hybrid Corn

Treatment of BOF + PA Doses	Plant Height (cm)	Number of Leaves (Sheet)	Cob Length (cm)	Cob Diameter (cm)	Weight 100 biji	Weight of Seed per Plot	Dry Weight of Crop Residue
T ₀	126,78	10,51	12,15	3,66	27,11	449,56	47,17
T ₁	133,56	10,71	12,89	3,77	27,78	479,33	52,25
T ₂	1358,80	10,82	13,79	3,96	27,44	482,89	62,01
T ₃	156,47	11,67	14,16	4,04	28,56	526,78	67,99
T ₄	138,02	11,29	12,94	3,89	28,11	508,78	64,11

Note: Description: T₀ = BOF + 0% N, P, K; T₁ = BOF + 25% N, P, K; T₂ = BOF + 50% N, P, K; T₃ = BOF + 75% N, P, K; T₄ = BOF + 100% N, P, K.

Table 4: Effect of Varieties of the Variables of Growth and Production of Hybrid Corn

Treatment of Corn Varieties	Plant Height (cm)	Number of Leaves (Sheet)	Cob Length (cm)	Cob Diameter (cm)	Weight of 100 Seeds	Weight of Seed per Plot	Dry Weight of Crop Residue (g)
V ₁	164,63 b	11,93b	14,39	4,20 b	28,67	591,00 b	73,74 b
V ₂	122,91 a	10,25a	12,05	3,55 a	27,27	416,40 a	44,94 a
V ₃	126,84 ab	10,81a	13,12	3,85 ab	27,47	461,00 a	57,45 a
HSD 0,05	39,95	1,69	ns	0,54	ns	112,15	21,45

Note: Description: The numbers followed by the same letters mean different non significant. V₁ = Pioneer 27; V₂ = Bisi 18; V₃ = DK 85.

Table 5: The Effect of the Interaction of Biological Organic Fertilizer Plus N, P, K and Varieties of the Variables of Growth and Production of Hybrid Corn



Treatment of BOF + Non Organic and Corn Varieties	Plant Height (cm)	Number of Leaves (Sheet)	Cob Length (cm)	Cob Diameter (cm)	Weight of 100 Seeds	Weight of Seed per Plot	Dry Weight of Crop Residue (g)
T ₀ V ₁	138,27	11,27	12,79	3,96	27,67	487,33	68,95
T ₀ V ₂	118,40	9,80	10,95	3,21	26,67	403,33	30,09
T ₀ V ₃	123,67	10,47	12,71	3,79	27,00	458,00	52,47
T ₁ V ₁	158,87	11,67	13,55	3,99	28,33	504,00	61,02
T ₁ V ₂	116,20	9,93	12,42	3,49	28,00	453,00	42,39
T ₁ V ₃	125,60	10,53	12,73	3,84	27,00	481,00	53,33
T ₂ V ₁	177,53	12,27	15,61	4,45	29,33	576,67	80,49
T ₂ V ₂	108,73	10,40	12,36	3,60	27,33	422,00	51,89
T ₂ V ₃	127,80	11,20	13,42	3,85	27,67	450,00	59,95
T ₃ V ₁	185,80	12,67	16,30	4,49	29,67	714,67	88,97
T ₃ V ₂	152,07	11,07	12,73	3,73	27,67	391,00	52,47
T ₃ V ₃	131,53	11,27	13,45	3,88	28,33	474,67	62,53
T ₄ V ₁	162,67	11,80	13,72	4,12	28,33	672,33	79,24
T ₄ V ₂	119,13	10,07	11,81	3,70	26,67	412,67	47,85
T ₄ V ₃	125,60	10,60	13,31	3,87	27,33	441,33	58,95

Note: Description: T₀V₁ = BOF + 0% N, P, K dan Pionner; T₀V₂ = BOF + 0% N, P, K dan BISI 18; T₀V₃ = BOF + 0% N, P, K dan DK 85; T₁V₁ = BOF + 25% N, P, K dan Pionner; T₁V₂ = BOF + 25% N, P, K dan BISI 18; T₁V₃ = BOF + 25% N, P, K dan DK 85; T₂V₁ = BOF + 50% N, P, K dan Pionner; T₂V₂ = BOF + 50% N, P, K dan BISI 18; T₂V₃ = BOF + 50% N, P, K dan DK 85; T₃V₁ = BOF + 75% N, P, K dan Pionner; T₃V₂ = BOF + 75% N, P, K dan BISI 18; T₃V₃ = BOF + 75% N, P, K dan DK 85; T₄V₁ = BOF + 100% N, P, K dan Pionner; T₄V₂ = BOF + 100% N, P, K dan BISI 18; T₄V₃ = BOF + 100% N, P, K dan DK 85.

MST) can be seen in Table 6. The average uptake of N, P and K from the three varieties of corn grown in the highest fertilizer treatment BOF + 75% N, P, K. The highest response to the uptake of N, P and K are varieties of Pioneer 27 amounted to 3.31% of N, 0.214% P and 2.90% K; Bisi 18 varieties of 3.00% N, 0.204%

P and 2.51% K; DK 85 varieties amounted to 2.64% of N, 0.185% P and 2.32% K.

Observations of weeds is done before planting and harvest. The results of the analysis of vegetation weeds before planting with the calculation of Summed Dominance Ratio (SDR), fresh weight and dry weight of weeds are listed in Table 7.



Weed Vegetation Analysis at 2 Days Before Harvest

The results of the analysis of weed vegetation 2 days before harvest showed that a decline in weed species. SDR results in all treatment plots show, that is the dominant weed *Cyperus compesus*. The results of the analysis of weed vegetation (SDR) are listed in Table 7.

DISCUSSION

Treatment without inorganic fertilizer (N, P, K) in this study showed the lowest value of both the variables of growth and yield components of maize, as also happens in the treatment of the interaction of BOF + N, P, K deliver growth and better results than without fertilizer inorganic.

Tabel 6: The Effect of Interaction Treatment on Nutrient Uptake NPK (%)

Treatment of BOF+ Non Organic Fertilizer	Varieties		
	V1	V2	V3
	Nutrient Uptake N%
T0	2,52	2,30	2,21
T1	2,54	2,31	2,55
T2	2,74	2,45	2,51
T3	3,31	3,00	2,64
T4	2,79	2,89	2,61
Treatment of BOF+ Non Organic Fertilizer	Varieties		
	V1	V2	V3
	Nutrient Uptake P%
T0	0,153	0,102	0,159
T1	0,147	0,135	0,171
T2	0,171	0,192	0,162
T3	0,214	0,204	0,185
T4	0,172	0,194	0,181
Treatment of BOF+ Non Organic Fertilizer	Varieties		
	V1	V2	V3
	Serapan Hara K%
T0	2,17	2,20	1,92
T1	2,41	2,13	2,02
T2	2,19	2,23	2,26
T3	2,90	2,51	2,32
T4	2,75	2,26	2,28

Tabel 7: Result of Weed Vegetation Analysis Before Land Tillage

No.	Type of Weed	SDR (%)	Wet Weight (g)	Dry Weight (g)
1	<i>Imperata cylindrica</i>	69,54	1492	566
2	<i>Cyperus rotundus</i>	8,72	22,0	6,0
3	<i>Borreria alata</i>	6,39	36,0	14,0
4	<i>Ageratum conyzoides</i>	5,46	206	7,9
5	<i>Micania micrantha</i>	3,39	68,0	26,5
6	<i>Melaleuca spp</i>	3,39	4,0	1,0
7	<i>Acrostichum aureum</i>	3,11	2,0	0,5

Table 8: Average of Weed Vegetation Analysis (SDR) 2 Days Before Harvest (%)

No	Treatment	Cyperus Compresus	Name of Weed	Borreria repens Dc.	Amount (%)
			Hyptis Capitata		
1	T0V1	80,88	11,12	8	100
2	T0V2	71,17	0	28,83	100
3	T0V3	66,55	34,45	0	100
4	T1V1	48,15	30,24	31,61	100
5	T1V2	79,36	10	10,64	100
6	T1V3	58,41	11,59	33	100
7	T2V1	44,3	42,47	13,23	100
8	T2V2	48,35	29,24	22,41	100
9	T2V3	26,05	33,33	40,62	100
10	T3V1	100	0	0	100
11	T3V2	65,5	34,5	0	100
12	T3V3	43,3	43,3	13,4	100
13	T4V1	26,3	73,7	0	100
14	T4V2	74,32	21,24	4,44	100
15	T4V3	93,07	6,93	0	100

Note: Description: T₀V₁ = BOF + 0% N, P, K with Pionner; T₀V₂ = BOF + 0% N, P, K with BISI 18; T₀V₃ = BOF + 0% N, P, K with DK 85; T₁V₁ = BOF + 25% N, P, K with Pionner; T₁V₂ = BOF + 25% N, P, K with BISI 18; T₁V₃ = BOF + 25% N, P, K with DK 85; T₂V₁ = BOF + 50% N, P, K with Pionner; T₂V₂ = BOF + 50% N, P, K with BISI 18; T₂V₃ = BOF + 50% N, P, K with DK 85; T₃V₁ = BOF + 75% N, P, K with Pionner; T₃V₂ = BOF + 75% N, P, K with BISI 18; T₃V₃ = BOF + 75% N, P, K with DK 85; T₄V₁ = BOF + 100% N, P, K with Pionner; T₄V₂ = BOF + 100% N, P, K with BISI 18; T₄V₃ = BOF + 100% N, P, K with DK 85.

Further enhanced the provision of inorganic fertilizer also increases the growth and yield components of maize at 75% dose recommended doses of inorganic fertilizer were tested. This shows that the administration of BOF alone in sub-optimal land is not enough to support growth and crop production, then added inorganic fertilizer to accelerate the availability of nutrients that plants need, because of the nature of organic fertilizers slow release in the supply of nutrients while inorganic fertilizer quickly available in a number of sufficient. In line proposed (Agustina,

1990; and Jumin, 2005) that the availability of nutrients in sufficient quantity and balance are the main factors that determine the success rate of growth and maximum production.

HSD test results, 27 Pioneers hybrid corn varieties produce the best growth and production compared Bisi 18 and DK 85, visible from all variables corn growth and yield components showed the highest values (Table 4). Suspected of hybrid corn variety Pioneer 27 has the adaptability of the environment in swampy areas are better than the varieties and varieties Bisi 18 DK 85. In line with the opinion of Toha (2008) in Barus *et al.* (2014), that the potential yield of a particular variety is not can be separated with the level of adaptation and stability in a growing environment. The level of production of a crop depending on the variety, cultivation methods and environmental conditions where the plants grow. Suistability a cultivated plants against the growth environment is very effect to growth and productivity of these plants. Furthermore, according Simatupang (1997) in Biological *et al.* (2011), that the increase in productivity due to a variety of these varieties able to adapt to the growth environment. While the hybrid maize varieties Bisi 18 have a genetic trait with poorly adapted to the environment swampy areas, so it is not able to demonstrate the primacy of nature, as a result of growth and production produced less than optimal. In line with the opinion of Lovelles (1989) in Biological *et al.* (2011), that each plant has a certain tolerance range of the environmental conditions. Therefore, most plants can be successfully grown in diverse conditions.

The interaction of biological treatment of organic fertilizer + N, P, K with some hybrid corn varieties in swampy areas no real effect on all observed variables. It is suspected between the

factors of fertilizer varieties not cooperate in influencing the growth and production of several varieties of hybrid corn plants or both factors influence treatment separately. According to Hanafi (2010), that is not the interaction of two factors of treatment due to two factors are not able to work together so that the mechanism of action is different or one of the factors did not play a role in an optimal or even antagonistic, which press against each other and their respective influence. Although statistically the interaction of both treatment factors are not real, but a tabulation seen the difference. Combination treatment of BOF + 75% N, P, K with 27 Pioneer hybrid corn varieties produce the best growth and production compared to the combination of other treatments (Table 5). Allegedly at BOF + 75% N, P, K is able to provide sufficient nutrients and balanced to support the growth, development and production of hybrid corn variety Pioneer 27. In line with the opinion of Biological *et al.* (2006) in Pasta *et al.* (2015) that growth, production and quality of sweet corn is influenced by genetic factors that variety and environmental factors such as soil fertility. There is a positive correlation between a combination of BOF + N, P, K, which is further enhanced with plant height, weight of crop residue and dry weight of seed per plot (Figure 1 and Figure 2). In the graph shows that the higher the dose of fertilizer given a high increase in corn and positively correlated with the weight of shelled corn kernels per plot to the extent of fertilizer combination BOF + 75% N, P, K. Similarly, what happened in Figure 2 shows that the increase in plant height due to the increase in dry weight of crop residue of corn and fertilizer peak in the combination treatment of BOF + 75% N, P, K. This is in line propounded by Sujana *et al.* (1991) that generally corn hybrids provide a higher yield than the open pollinated

maize. The study showed a positive correlation between the corn is tall with a yield of corn.

Besides the presence of weeds fertilizer factor in corn planting very effect on the growth and production of maize. Generally weeds that predominate in the area of the corn crop that is kind of puzzle-tekian as found in this study there are a lot of weeds *Cyperus compressus*. This includes family Cyperaceae weeds that have a high ability to adapt to the environment, multiply by seed and rhizome, can grow in extreme environmental conditions. As a result of this weed can be controlled space and strong growth in competing with cultivated plants. In accordance opinions Holm *et al.* (1988) in Suryaningsih (2011), that the family Cyperaceae weeds, including weeds that have high adaptability to the environment and a strong root rhizomes and can reproduce by seed and rhizome. Furthermore, according to Tjitrosoedirjo (2010), that the family Cyperaceae weeds have long root systems, many have seeds that cause rapid spread.

The results of the analysis of vegetation weeds before planting No. 7 of weed species (Table 6) and that the dominant weed *Imperata Cylindrica* (69.54%). After treatment, there are 3 types of weeds (Table 7) but the dominant weed namely *Cyperus compressus* (100%), on average the highest dominant weed in the combination treatment dose interaction BOF + 75% N, P, K. This suggests that the response of weeds and corn plants against fertilizer is the same in terms of fighting nutrients and other growth factors. Besides, the use of Biological Organic Fertilizer (BOF) are weed seeds that are still dormant so it will grow on the environmental conditions favorable. Along expressed Fitriana *et al.* (2011) that the use of manure can lead to the development of weeds in cultivated land.

Physical condition swampy areas that are less supportive and nutrient-poor as well as the presence of weeds although the application of organic and inorganic fertilizers have not been sufficient to support plant growth due to maintenance should still be maintained. In line with the opinion of Suryaningsih (2011) that the presence of weeds can harm other plants that are nearby or a staple crop. Corn is very sensitive to weed competition with the decrease in the yield of 16-56% (Violic, 2000). Critical period of maize crops to compete with weeds occurred on days 20 and 45, then the critical period of maize plants also occurred on days 80 to 150 (Sembodo, 2010). Physically weeds compete with plants in the utilization of space, light and chemically in the utilization of water, nutrients, gases are important in the process allelopati. Competition can take place when components or substances needed by weeds or crops that are in limited quantities, and closely spaced together required (Moenandir, 2010).

CONCLUSION

1. Treatment of organic biological fertilizer + 75% N, P, K is the best treatment for the growth and production of hybrid corn in swampy areas.
2. The hybrid corn varieties are varieties that are best Pioneer 27 the growth and production in swampy areas with yield per plot of 714.67 kg (3, 57 ton ha).
3. Weeds dominant before the research is *Imperata cylindrica* and the in harvesting is *Cyperus rotundus*.

ACKNOWLEDGMENT

Author to convey to the Ministry of Research, Technology and Higher Education Directorate

General for Strengthening Research and Development that has funded these activities through Budget Competitive Research at Year 1 Budget Year 2016.

REFERENCES

1. Agustina (1990), *Nutrisi Tanaman*, Rineka Cipta, Jakarta.
2. Alihamsyah T (2005), "Pengembangan Lahan Rawa Lebak Mitra Usaha Pertanian", Balittra, Banjarbaru, 53 Hal.
3. Ardjasa W S dan and Bangun P (1985), "Pengendalian Gulma Pada Tanaman Pokok", Pusat Penelitian dan Pengembangan Pangan, Bogor.
4. Balai Penelitian Tanah (2005), "Petunjuk Teknis Analisis Kimia Tanah, Tanaman, Air dan Pupuk", Badan Penelitian dan Pengembangan Pertanian, Departemen Pertanian.
5. Barus W A, Khair dan H and Siregar M A (2014), "Respon Pertumbuhan dan Produksi Kacang Hijau (*Phaseolus radiatus* L.) Akibat Penggunaan Pupuk Organik Cair dan Pupuk TSP", *Journal Agrium*, Vol. 19, No. 1.
6. Badan Pusat Statistik (2012), "Indikator Pembangunan Provinsi Sumatera Selatan Tahun 2012", <http://sumsel.bps.go.id> [18 April 2012].
7. Balai Pengkajian Teknologi Pertanian Sumatra Selatan (2001), "Peta Zona Agro Ekologi Propinsi Sumatra Selatan", BPTP Sumsel, Palembang.
8. Djafar Z R, Syopyan J, Dartius A, Zainal D, Sunyati E and Hadiono dan Sagiman (1990), "Dasar-Dasar Agronomi", BKS-B USAID, Palembang.

9. Fitriana M, Parto Y, Munandar dan and Budianta D (2013), "Pergeseran Jenis Gulma Akibat Perlakuan Bahan Organik pada Lahan Kering Bekas Tanaman JAgung (*Zea mays* L.)", *Jurnal Agron Indonesia*, Vol. 41, No. 2, pp. 118-125.
10. Gardjito M, Djuwardi A and Harmayani E (2013), "Pangan Nusantara. Karakteristik dan prospek Untuk Percepatan Diversifikasi pangan", Penerbit Kencana Prenada Media Group, Edisi pertama. Jakarta.
11. Hanafiah K A (2010), "Rancangan Percobaan", Rajawali Pers, Jakarta.
12. Hanafiah K A (2005), "Dasar-dasar Ilmu Tanah", RajaGrafindo, Jakarta Hardjowigeno S, 2010, Ilmu Tanah, Akademika Pressindo, Jakarta.
13. Hayati M and Hayati dan Nurfandi E (2011), "Pengaruh Pupuk Organik dan Anorganik Terhadap Pertumbuhan Beberapa Varietas Jagung Manis Dilahan Tsunami", *Jurnal Floratek*, Vol. 6, pp. 74-83.
14. Jumini Nurhayati dan Murzani (2011), "Efek Kombinasi Dosis Pupuk N, P, K dan Cara Pemupukan Terhadap Pertumbuhan dan Hasil Jagung Manis", *Jurnal Floratek*, Vol. 6, pp. 165-170.
15. Khair H, Pasaribu dan M S and Suprpto E (2013), "Respon Pertumbuhan dan Produksi Tanaman Jagung (*Zea mays* L.) Terhadap Pemberian Pupuk Kandang Ayam dan Pupuk Organik Cair Plus", *Jurnal Agrium*, Vol. 18, No. 1.
16. Kresnatita S, Koesriharti dan and Santoso M (2004), "Pengaruh Pemberian Pupuk Organik dan Nitrogen Terhadap Pertumbuhan dan Hasil Tanaman Jagung Manis", Fakultas Pertanian, Universitas Brawijaya.
17. Lakitan B (2007), "Fisiologi Tumbuhan dan Perkembangan Tanaman", Raja Grafindo, Jakarta.
18. Mahdil S and Hassan G I (2010), "Biofertilizersin Organic Agriculture", *Phytology*, Vol. 2, No. 10, pp. 42-45.
19. Merlin *et al.* (2011), "Pusat Penelitian Tanah 1983", Terms of Reference Type. As. "P3TT Bogor, Mahdiannoor Efektivitas Pemberian Trichoderma Spp. dan Dosis Pupuk Kandang Kotoran Ayam pada Lahan Rawa Lebak Terhadap Pertumbuhan dan Hasil Tanaman Kacang Panjang (*Vigna Sinensis* L.)", *Jurnal Ziraa'ah*, Vol. 33, No. 1, pp. 91-98.
20. Murbandono H S (1998), "Membuat Kompos", Penebar Swadaya, Jakarta.
21. Marschner H (1989), "Mineral Nutrition of Higher Plant", Academic Press, London.
22. Moenandir J (2010), "Ilmu Gulma", Universitas Brawijaya Press, Malang.
23. Pangaribuan D H, Yasir dan M and Utami N K (2012), "Dampak Bokashi Kotoran Ternak dalam Pengurangan Pemakaian Pupuk Anorganik pada Budidaya Tanaman Tomat", *Jurnal Agron Indonesia*, Vol. 40, No. 3, pp. 204-210.
24. Pasta I, Ette dan A and Barus H N (2015), "Tanggap Pertumbuhan dan Hasil Tanaman Jagung Manis (*Zea mays* Saccharata Sturt) pada Aplikasi Berbagai Pupuk Organik", *Jurnal Agrotekbis*, Vol. 3, No. 2, pp. 160-177.
25. Pusat Statistik (2012), "Indikator Pembangunan Provinsi Sumatera Selatan

- Tahun 2012”, <http://sumsel.bps.go.id> [18 April 2012].
26. Risnandar C (2011), “Jenis dan karakteristik pupuk kandang”, <http://alamatani.com/pupuk-kandang.html>, di akses april 2015.
 27. Rismunandar (1992), “Tanah dan Sifat-Sifansya Bagi Pertanian”, Sinar Baru, Bandung.
 28. Rukmana R (1995), “Usaha Tani Jagung”, Kanisius, Yogyakarta.
 29. Setiawan A E (2009), “Memfaatkan Kotoran Ternak”, Penebar Swadaya, Jakarta.
 30. Sastrosoedirdjo *et al.* (2010).
 31. Subagyo A (2006), “Lahan Rawa Lebak dalam Didi Ardis S *et al.* Karakteristik dan Pengelolaan Lahan Rawa”, Balai besar Penelitian dan Pengembangan Sumber Daya Lahan Pertanian, Bogor, Hal, pp. 99-116.
 32. Suryaningsih M and Joni A A K Darmadi (2011), “Inventarisasi Gulma pada Tanaman Jagung (*Zea mays* L.) dilahan Sawah Kelurahan Padang Galak Denpasar Timur Kodya Denpasar”, *Provinsi Bali. Jurnal Simbiosis*, Vol. 1, No. 1, pp. 1-8.
 33. Sutanto R (2002), “Pertanian Organik Menuju Pertanian Alternatif dan Berkelanjutan”, Kanisius, Yogyakarta.
 34. Sutedjo M M (2008), “Pupuk dan Cara Pemupukan”, Rineka Cipta, Jakarta.
 35. Syafriani E, Sodikin E and Priadi dan Munandar D P (2004), “Pengaruh Pupuk Hayati pada Berbagai Variasi Dosis Pupuk Kandang dan Pupuk NPK Terhadap Pertumbuhan dan Produksi Tanaman Jagung Varietas C-7”, *Jurnal Tanaman Tropika*, Vol. 7, No. 2, pp. 69-77.
 36. Sembodo D R J (2010), “Gulma dan Pengolahannya”, Penerbit Graha Ilmu, Edisi Pertama, Yogyakarta.
 37. Siburian R (2007), “Pengaruh Konsentrasi dan Waktu Inkubasi EM4 terhadap Kualitas Kimia Kompos”, Fakultas Sains dan Teknik, Universitas Nusa Cendana, Kupang.
 38. Simamora (2006), “Pengaruh Waktu Penyiangan dan Jarak Tanam terhadap Pertumbuhan dan Produksi Tanaman Jagung (*Zea Mays* L) Varietas DK 3”, <http://Repository.Usu.Ac.Id/123456789/7568/1/09E00237.Pdf>, Diakses2 April 2015.
 39. Tufaila M, Laksana dan D D and Alam S (2014), “Aplikasi Kompos Kotoran Ayam untuk Meningkatkan Hasil Tanaman Mentimun (*Cucumis sativus* L.) pada Tanah Masam”, *Jurnal Agroteknos*, Vol. 4, No. 2, pp. 119-126.
 40. Warisno (2009), “Tanaman Jagung Hibrida”, <http://Www.Digilib.Uns.Ac.Id/Upload/Dokumen/173072312201007154>, Pdf, DiaksesMaret 2015.
 41. Wijaya K A (2008), “Nutrisi Tanaman”, Prestasi Pustaka, Jakarta.
 42. Yunus M (1991), “Pengelolaan Limbah Peternakan”, Jurusan Produksi Ternak LUW-Universitas Brawijaya, Animal Husbandry Project.



International Journal of Engineering Research and Science & Technology

Hyderabad, INDIA. Ph: +91-09441351700, 09059645577

E-mail: editorijerst@gmail.com or editor@ijerst.com

Website: www.ijerst.com

