



International Journal of Engineering Research and Science & Technology

ISSN : 2319-5991
Vol. 6, No. 1
February 2017



www.ijerst.com

Email: editorijerst@gmail.com or editor@ijerst.com

Research Paper

AGRONOMICAL CHARACTERISTICS PERFORMANCE OF F₁ GENERATION RAINFED RICE FROM CROSS BREEDING BETWEEN INTRODUCED RICE VARIETY AND LOCAL RICE VARIETY OF SOUTH SUMATRA

Gusmiatun^{1*}

*Corresponding Author: Gusmiatun ✉ gusmiatun69@yahoo.com

The effort to improve performance and quality of rainfed rice currently can be done by using cross breeding between local rice variety and introduced rice variety. This local rice variety had aromatic gene with specific aroma as well as taste and texture which was in accordance to community cultural pattern of South Sumatra. This aromatic gen was found in Dayang Rindu rice variety, whereas the introduced rice variety had advantages in term of age, production and adaptation capability in South Sumatra area. The research objective was to determine the performance of F₁ generation crop resulting from cross breeding between introduced rice variety and local rice variety indigenous of South Sumatra in order to produce new superior rice variety having characteristics of high production, short age as well as specific taste and aroma of South Sumatra specialty. This research was conducted at farm plot in Pulau Semambu Village, Indralaya Utara Subdistrict from January to April 2016. The method used in this research was Randomized Block Design with one treatment factor and six replications for each treatment. Rice seeds used as treatments were consisted of Inpago-7, JatiLuhur, Dayang Rindu and F₁ generation varieties. The results showed that F₁ generation rice variety had no significant different in term of crop height and tiller numbers than Jati Luhur brood which had high yield potential (3.5 to 4 ton/ha) and crop's age of 110 to 118 days so that it had potential to be developed as candidate of new superior rice variety.

Keywords: Agronomical characteristics, Rainfed rice, F₁ generation

INTRODUCTION

The food autonomy at province level should be achieved in order to create national food tenacity.

One of strategies that can be implemented is by utilizing unproductive land which consisted of dry land and swamp land for rice cultivation. South

¹ Muhammadiyah University, Palembang 30263.

Sumatra Province had extensive dry land area in which only 52,679 ha that had been managed or about 13.7% from total dry land area of 385,407 ha (BPTP Sumatera Selatan, 2009).

The opportunity of rainfed rice development at dry land in addition to traditional land as second crop was in accordance to new land clearing for plantation area. The second crop can be cultivated up to third year at young rubber plantation area as well as up to fourth year at oil palm plantation area (Suryana, 2008; and Yusuf, 2009). Rainfed rice can also be cultivated at swamp land area during dry season (Gusmiatun, 2015). Although rainfed rice has high development potential, but farmers eagerness to cultivate rainfed rice was relatively low. This was caused by low productivity and low quality of local variety rainfed rice which was generally used by farmers. The average production of this local variety rainfed rice was in the range of 1.5 to 2.5 ton/ha or about 43% from productivity of paddy field rice (BPS, 2007).

Despite its mentioned diadvantages, there was local variety rainfed rice which had specific taste and wangi aroma that only available in certain area. Local variety rainfed rice which had specific taste, fragrant aroma, good tasting as well high economic value known as Dayang Putri was found in Musi Rawas District of South Sumatra Province. It is preferred by local community members. The disadvantages of Dayang Putri rice variety were long age of six months and low productivity with magnitude of 1.9 to 2.0 ton/ha (Portal Nasional RI, 2012). If rice having superior characteristics (aroma and taste) is cross bred with rice variety having short age and high production, then it is expected to yield the new rice variety having short age and high production and fragrant aroma which in turn can increase the interest of farmers to cultivate rainfed rice.

The government up to nowadays had been introduced several superior rice varieties, but the assembly of new superior rice varieties by considering the interest of region in which the varieties will be developed is still needed because the community preferences toward taste are different amongst areas/regions. The objective of this effort is that farmers are eager to adopt new rice varieties. Results of field survey showed that most farmers at South Sumatra are still cultivating local rice varieties because this local rice varieties have taste in accordance to farmers community preferences.

The assembly of new variety is started from cross breeding between the selected brood, crop variation from the cross breeding results of F_1 generation is depends on brood selection which will produce heterotic hybrid. Hybrid vigor heterosis is a trend in which F_1 individual from cross breeding results will have better performance than one of brood or average of both broods.

For self pollination crop, the probability of heterosis utilization is started with brood selection which produce the best characteristics combination. This is important to proceed breeding for the best gene combination that can be obtained from variety that is relatively homozygote (Hayes, 1964).

The advantages of unhulled rice potential should be determined for F_1 hybrids from cross breeding between local rice variety and superior rice variety in order to produce new hybrid rice having high production and fragrant aroma. This research objective was to determine the growth and production of F_1 crop resulting from cross breeding between Jati Luhur rice variety and local rice variety of Dayang Rindu.

MATERIALS AND METHOD

This research was conducted at experimental plots of Agricultural Faculty, Muhammadiyah University Palembang located in Pulau Semambu Village, Indralaya Utara Subdistrict from January to April 2016. Rice seeds used in this study were originated from four rice varieties consisting of broods of Jati Luhur, Dayang Rindu, F₁ (JTL-DR), and control variety of Inpago-7. Field research was conducted by using Randomized Block Design with one treatment factor and six replications for each treatment. Treatment unit was plot having size of 2 m x 5 m, two seeds was planted in a hole with planting distance of 25 cm x 25 cm.

Observation was consisted of crop height (cm), productive tiller numbers, harvest date (Day After Planting), percent of empty unhulled rice (%) and unhulled rice weight per clump (g).

RESULTS AND DISCUSSION

Results

1. Vegetative parameters (crop height, productive tiller numbers).
2. Production parameters (percent of empty unhulled rice, unhulled rice weight/clump).

No	Variety Crop Morphology	Jati Luhur (JTL)	Dayang Rindu (DR)	F ₁ (JTLxDR)	Inpago-7
		1	Crop height (cm)	105 a	137 c
2	Productive tiller numbers	13 a	9 b	12:00 AM	13 a
3	Percent of empty unhulled rice (%)	8.1 a	7.1 a	7.9 a	7.5 a
4	Unhulled rice weight/clump (g)	31.60b	28.15 a	31.50b	30.20b
5	Harvest age (Day After Planting = DAP)	95.50 a	142.75 c	95.75 a	114.00 b

Figure 1: (a) Height Differences and (b) Productive Tiller Numbers Differences of Rainfed Rice Crop Amongst Broods and F₁




Figure 2: (a) Differences of Empty Unhulled Rice Percentage and (b) Differences of Unhulled Rice Weight/Clump of Rainfed Rice Crop Amongst Broods and F₁





Figure 2 (Cont.)



3. Harvest age (DAP).

Figure 3: Differences of Rainfed Rice Age (DAP) Among Stbroods and F₁ Generation




Figure 4: F₁ (Jati Luhur X Dayang Rindu)




Figure 5: Inpago-7




Figure 6: Dayang Rindu





Figure 7: Jati Luhur



Discussion

The average crop height of F_1 generation produced from cross breeding of Jati Luhur rice variety and local rice variety of Dayang Rindu was 103 cm which was not significantly different than Jati Luhur variety height of 105 cm, but it was lower than Dayang Rindu variety height of 137 cm (Figure 1a). This results was the expected performance because higher crop is not always better than shorter crop, even higher crop is easily fall down especially during period of panicle filling. On the other hand, short and stiff stem is the expected characteristics for development of superior rice varieties because it is not only fall resistant, but also has more balance ratio of unhulled rice and rice straw (Jennings *et al.*, 1979; and Yoshida, 1981). The characteristic of hybrid crop (in this case is F_1 generation is determined by characteristic of its broods. If characteristics of both broods are combined synergically, then the produced generation will pose combine characteristics which is better than both of its brood (You *et al.*, 2006).

Tiller numbers produced by each variety are determine by crop's genetical factors. Results of the study showed that F_1 generation inherited brood gen of Jati Luhur which control numbers of productive tiller so that productive tiller numbers produced by F_1 generation was not different than productive tiller numbers produced by its brood of Jati Luhur with magnitude of 12 tillers. This magnitude was not different than tiller numbers produced by control variety of Inpago-7, whereas Dayang Rindu brood had only produced 9 tillers (Figure 1b). Capacity of tillers is one important characteristics for superior variety. Crops having high tiller numbers are suitable for several planting distance variations, capable to compensate the death clumps and capable to achieve leave area quickly (Yoshida, 1981).

The crop growth in term of crop height and productive tiller numbers affects the yield of unhulled rice. The higher the numbers of productive tiller, the higher the crop production. The F_1 generation had produced higher unhulled rice weight per clump (30.2 to 31.6 g/clump) than Dayang Rindu variety as shown in Figure 2a.

The effect of crop height on unhulled rice yield was found on Dayang Rindu rice variety which had the highest crop height so that it produced the lowest unhulled rice yield. According to Navasero and Tanaka (1966), stem elongation could results in assimilates competition with panicle development because stem elongation can consume about 60% of produced net assimilates and the rest of assimilates are distributed for leaves growth and panicle (Wada, 1969). Furthermore, Murata and Matsushima (1978) had stated that excessive vegetative growth can results in decrease of assimilates supply.

Research results for all tested crops (Jati Luhur, Dayang Rindu, F_1 and Inpago-7 varieties) showed low percentage of empty unhulled rice in the range of 7.1 to 8.1%. This finding showed that capability of *Sourceto* supply assimilates into Sink was relatively sufficient. The opposite condition was found on crops that had higher percentage of empty unhulled rice (more than 20%) because source was less capable to supply assimilates although it had high sink especially on unfavourable environment so that many sink was empty or not be utilized by source (Makarim *et al.*, 2004).

Harvest age of F_1 crop was not different than Jati Luhur brood (116 to 118 DAP), but its harvest umur was shorter than that of local tetua of Dayang Rindu with magnitude of 150 DAP (Figure 3). This finding was the expected generation from

both broods used in this study because one of characteristics for superior rice variety is short umur. The age of rice crop is determined by the period length of vegetative phase consisting growth phase of crop vegetative organs such as stem elongation or crop height De Datta (1981) and Yoshida (1981). The higher the rice crop height, the longer the vegetative phase period so that its harvest umur is longer. Local rice variety of Dayang Rindu had the highest crop height so that its harvest age is the longest.

CONCLUSION

1. The F_1 crop produced from cross breeding between Jati Luhur and Dayang Rindu varieties had similar growth and production levels with parent Jati Luhur. This crop height was 101 cm and unhulled rice production per clump was 31.5 g.
2. The F_1 crop had short age with magnitude of 118 Day After Planting.

REFERENCES

1. Council of Agricultural Technology Study of South Sumatra (2009), "Agroecological Zone Map of South Sumatra Province", BPTP of South Sumatra, Palembang.
2. Council of Statistical Center (2010), "Banyuasin in Numbers", Gowen (Ed.), pp. 49-65, Hafner Publishing Company, New York.
3. Gusmiatun *et al.* (2015), "Performance of Agronomical Characteristics of Rainfed Rice Varieties at Oganllir District, South Sumatra Province", *International Journal of Engineering Research and Science & Technology*, Vol. 5, No. 2, pp. 27-35, ISSN: 2319-5991.
4. Hayes H K (1964), "Development of the Heterosis Concept", in Heterosis J W, Jennings P R *et al.* (1979), "Rice Improvement", IRRI, Los Banos, Philippines.
5. Makarim A K (2004), "New Type of Rice: Cultivation by Using Integrated Crop Management Approach", *Rice Crop Research Council*, p. 48, ISBN: 979-540-021-5.
6. Navasero S A and Tanaka A (1966), "Low Light Induced Death of Lower Leaves of Rice and Its Effect on Grain Yield", *Plant and Soil*, Vol. 14, pp. 17-31.
7. Suryana A (2008), "Field Technical Guidance for Integrated Crop Management (ICM) of Rainfed Rice", *Agricultural Research and Development Council*, p. 7, Agricultural Department.
8. Wada G (1969), "The Effects of Nitrogenous Nutrition on the Yield Determining Process of Rice Plant", *Bull. Nat. Inst. Sci. Japan Ser. A*, Vol. 16, pp. 27-167.
9. You A, Lu X, Jin H, Ren X, Liu K, Yang G, Yang H, Zhu L and He G (2006), "Identification of Quantitative Trait Loci Across Recombinant Inbred Lines and Testcross Populations for Traits of Agronomic Importance in Rice", *Genetics*, Vol. 172, pp. 1287-1300.
10. Yoshida S (1981), *Fundamentals of Rice Crop Science*, International Rice Research Institute, Los Banos, Philippines.



International Journal of Engineering Research and Science & Technology

Hyderabad, INDIA. Ph: +91-09441351700, 09059645577

E-mail: editorijerst@gmail.com or editor@ijerst.com

Website: www.ijerst.com

