



International Journal of Engineering Research and Science & Technology

ISSN : 2319-5991
Vol. 6, No. 2
May 2017



www.ijerst.com

Email: editorijerst@gmail.com or editor@ijerst.com

Research Paper

EVALUATION OF SOME SOIL CHEMICAL PROPERTIES OF TIDAL SWAMP LAND AFTER LONG-TERM CULTIVATION

Dedik Budianta^{1*}, Ermatita², Napoleon¹, Agus Hermawan¹ and Harry Wijayanti³*Corresponding Author: Dedik Budianta ✉ dedik_budianto@yahoo.com

Tidal land is an acidic soil and has low fertility due to high acidity and low nutrients. The objective of this research is to evaluate the tidal land after long-term cultivation. Twelve soil samples of arable layers were collected at four typologies which are A, B, C and D types. Soil variables observed were pH, total N, organic C, Cation Exchange Capacity, P, Al and H-exchangeable and basic cations. After long-term cultivation, these soils had still high organic carbon and cation exchange capacity but low pH, Ca, Mg and total N. Relatively, high content of nutrient was found on Na and P. To improve infertility, farmers applied NPK in high rate which were 1000 kg ha⁻¹ for corn and 600 kg ha⁻¹ for rice. This soils had low pH and low nutrients. However these soil has potential to be cultivated for agriculture after improving pH and alleviating Al and pyrite.

Keywords: Tidal land, Acid sulphate soil, Macro elements, Pyrite, Land typology

INTRODUCTION

About 16.68 million ha mineral soils of tidal swamp land are distributed in Indonesia and from which about 4.19 million ha of acid potential soils have been reclaimed for agriculture and transmigration area and other of 5.34 million ha is still being used as agricultural development (Budianta and Windusari, 2016). Tidal swamp development in South Sumatra has been performed by Indonesian Government since 1969 which was started in PELITA I (The first five-year

development phase) 1969-1974 period that Indonesian Government had opened the tidal swamps about 22,000 ha in the province of South Sumatra and Jambi (Koswara and Rumawas, 1984), but Bugis farmers from South Sulawesi had previously developed this area since 1930s (Imanudin *et al.*, 2010). Tidal swamp areas in South Sumatra are located along eastern coast of Sumatra and approximately covered 2.92 million ha (Euroconsult, 1995). This tidal swamp is usually used for rice growth because in wet

¹ Department of Soil Science, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, 30662, South Sumatra, Indonesia.

² Department of Information System, Faculty of Computer Science, Universitas Sriwijaya, Indralaya, 30662, South Sumatra, Indonesia.

³ Department of Biology, Faculty of Mathematic and Natural Sciences, Universitas Sriwijaya, Indralaya, 30662, South Sumatra, Indonesia.


condition pyrite will be stable (Hairani *et al.*, 2005), but the most important constraint to rice growth on these soils were: (1) acidity (which includes the combined effect of pH, Al toxicity and P deficiency), and (2) Fe stress which was due to the combined effect of Fe toxicity and deficiency of other divalent cations such as Ca, Mg (Moore *et al.*, 1990). The total area of tidal swamp in South Sumatra that had been reclaimed is 373,000 ha and now this land is highly dependent on water supply from rainfall because high tide irrigation can not reach in the land (Imanudin *et al.*, 2010). The tidal swamp soil is characterized by poor chemical properties such as low soil pH and high in Fe and Al concentration (Masulili *et al.*, 2016) in combination with high rainfall and effect tidal movement make this area have excessive water in a certain period in the year (Purnomo *et al.*, 2008), especially in areas which have not been reclaimed. In these soils called acid sulfate soil (Dent, 1986), acid sulfate soil is formed in marine or brackish environment (Ilona *et al.*, 2012). Elements such as iron and aluminum are released to soil solution in large amount, while essential macro nutrients like phosphorus and nitrogen become fixed in the soil resulting in unavailable for plant growth (Dublin-Green and Ojanuga, 1988).

Acid sulfate soils with pH below 3.5 will have toxicities of H^+ , Al^{3+} , SO_4^{2-} and Fe^{3+} causing a decrease in soil fertility due to low soil bases and essential macronutrients such as N, P, K, Ca, Mg (Hairani *et al.*, 2005; Ristiani, 2012; and Budianta *et al.*, 2012) with values of 0.18% N, 14 mg kg^{-1} P, $0.40 \text{ (cmol(+))kg}^{-1}$ K, $1.08 \text{ (cmol(+))kg}^{-1}$ Ca and $1.30 \text{ (cmol(+))kg}^{-1}$ Mg (Nursanti, 2014) and low essential micro element such as Mn, Zn, Cu and Mo (Fami and Hanudin, 2008). Due to low fertility faced in acid sulfate soils, food crops cannot grow

optimal in these condition with low production (Hairani *et al.*, 2005), farmers should apply fertilizers in high rate. In this infertile soil, food crop will be produced in low quantity. Thus, the objectives of this study is to evaluate some soil chemical properties of tidal swamp land that has been cultivated more than 30 years for food crops (corn and rice) production in South Sumatra, Indonesia.

MATERIALS AND METHODS

Some soil samples were collected at arable layer from different tidal swamp typologies which are A, B, C, and D respectively. All soil samples were located in Banyuasin Regency covered about 2,489.55 ha and distributed in Mulia Sari and Banyuurip Villages belonged to Telang II and divided into 12 soil samples (Figure 1). Geographically soil samples were taken in coordinates locations about S $02^{\circ}41'24''$, E $104^{\circ}45'25''$ for A, S $02^{\circ}39'26''$ E $104^{\circ}44'27''$ for B, S $02^{\circ}39'44''$, E $104^{\circ}44'027''$ for C, and S $02^{\circ}38'37''$, E



104°43'16" for D land typologies respectively. All soil samples were taken in dry season and the A land typology condition was bare soil with some swamp grasses growth, meanwhile in B, C, and D typologies respectively were planted corn in mature condition. Farmers always fertilizes the rice with dosage of 600 kg NPK per ha and 1, 0 t ha⁻¹ for corn crops.

The soil samples were then brought to the laboratory of Soil Science Department, Faculty of Agriculture, Universitas Sriwijaya to prepare for soil analysis. The samples were dried, ground and sieved with 0.5 mm sieve diameter. Variables observed were soil pH (1;1, soil:water (Tohmas, 1996), organic-carbon (Nelson and Sommers, 1996), total N (Bremner, 1996), P (Bray-I) (Kuo, 1996), K, Na, Ca and Mg-exchangeability (Ammonium acetate pH: 7.0 (Helmke and Sparks, 1996; and Suarez, 1996). Na and K determined by a flame photometer and Ca and Mg were determined by atomic absorption spectro photometer. Cation Exchange Capacity (CEC) was determined by extraction using the NH₄Oac pH:7 (Sumner and Miller, 1996) and Al-exchangeability were extracted by 1 N KCl (Bertsch and Bloom, 1996). Pyrite was determined by measurement of Sulfur (Tabatabai, 1996).

RESULTS DAN DISCUSSION

Landuse Condition at Soil Sampling Areas

Twelve soil samples of arable layer at the depth of 0-20 cm which were distributed in four distinguished typologies were collected from farming areas in the dry season, in August 2016. The sampling were taken from two transmigran areas of Telang II of Tanjung Lago, Banyuasin Regency, South Sumatra, Indonesia and located

in Mulia Sari and Banyu Urip villages. Since 1981, this tidal swamp had been reclaimed and about 13,800 ha of the areas are potentially for agriculture (Marlina *et al.*, 2015). In the two places of soil samples consists of 4 different land typologies of swampy land which are A and B typologies located in Mulia Sari and C and D typologies were placed in Banyu Urip. The agricultural landuse in the vicinity of research area can be seen in Table 1.

The four types of swampy tidal land typologies in Mulia Sari and Banyu Urip based on the overflow of water consisted of (1) type A, flooded both during low and high level of tide, (2) type B, flooded only when high level of tide, (3) type C, never flooded even during high level of tide and the depth of ground water < 50 cm, and (4) type D, never flooded even during level high of tide and the depth

Table 1: Distribution of Agricultural Landuse in the Vicinity of Research Area (Intimulya Multikenncana, 2009)

No	Villages	Agricultural Coverage (ha)	Type of Crop
1	Tegal Sari	1,787.27	Coconut: 48.17 ha
			Oil Palm: 135.43 ha
			Ric : 1,603.40 ha
			Tertier blok average: 44 ha
2	Mulia Sari	1,057.70	All rice crop Tertier blok average: 44 ha
3	Banyu Urip	1,431.85	All rice crop Tertierblok average: 44 ha
4	Bangun Sari	1,431.85	All rice crop Tertier blok average: 44 ha
5	Sumber Mekar	785.93	All rice crop Tertier blok avergae: 44 ha

of ground water > 50 cm (Widjaja Adhi *et al.*, 1992; and Rahim *et al.*, 2015). During soil sampling at the two villages, conducted in dry season and in Mulia Sari, were found type A and B in south region at the nineteenth of south bridge and third of north bridge respectively (Rahmi *et al.*, 2015), meanwhile type C and D were obtained in Banyu Urip at the fourth of south bridge and at the fourth of north bridge respectively. With high rate fertilization in these Swamp lands, it was harvested around 4-7.5 t/ha rice in Mulia Sari. Meanwhile, type C of the swamp land in South Kalimantan was planted soybean, after the soil improved by liming and fertility and getting increase of soybean about 2.4 t/ha⁻¹ (Wijanarko *et al.*, 2016).

Evaluation of Tidal Swamp Land After Longterm Cultivation

Tidal swamp lands are generally considered as a marginal and fragile ecosystems, but the lands have potential to be utilized for the development of food crops such as rice, soybean and corn, as well as horticulture crops like citrus and vegetables (Koesrini *et al.*, 2013; and Koesrini *et al.*, 2014). According to Widjaya Adhi *et al.* (1992), that wetlands are distinguished based on arrival, influences the tidal water in the rainy season and the influence of sea water in the dry season, is divided into three zones, namely: 1) tidal brackish (zone I), 2) tidal fresh water (zone II) and non tidal (zone III). The tidal land based on agro-ecosystem can be divided into four main typologies of potential lands, acid sulfate lands, peat lands and saline lands. 1) Potential land is a land of upper layer of 0-50 cm, has a low pyrite content of 2% and has not undergone a process of oxidation. 2) Acid sulfate land is a land that has a layer of pyrite or sulfidic at depth <50 cm and all the land that has sulfuric layer, although the

depth of the layer pyrite > 50 cm. Pyrite layer is a layer or layers of sulfidic soil with the pyrite levels > 2%. Sulfuric horizon is a layer that shows the jerosite (brownlayer) or pyrite oxidation process pH (H₂O) <3.5. Acid sulfate land is divided into (i) actual acid sulfate land, and (ii) potential acid sulfate land that is not or has not undergone the process of pyrite oxidation. 3) Peatlands are wetlands that have a layer of peat and peat thickness is characterized by shallow peat (thickness 50-100 cm), peat medium (thickness of 100-200 cm), deep peat (200-300 cm) and deep peat (>300 cm). Mukhtar and Adiprasetyo (1993) said that peats have great prospects for the cultivation of crops. For coconut and oil palm cultivation can be carried out on peat medium and deep. 4) Saline land is land that received saline brines influence, when under the influence of seawater/salty over 4 months of the year and the content of Na in soil solution of 8% to 15%. Tidal swampland based hydrotopography divided into four types which require different management. Type A is the area of the swamp which is always a big tide overflow and small pairs. Type B is the only land with a tidal overflow. Type C is a land that is not overflowing with the tide, both pairs of large and small pairs, but the depth of ground water is less than 50 cm from the soil surface. Type D does not overflow tidal land both pairs of large and small pairs, but the depth of ground water for more than 50 cm from the soil surface. Based on agro-ecosystem, Mulia Sari and Banyu Urip have potential acid sulfate lands and tidal fresh water. Farmers plant rice, corn and coconut in their land to fulfill their life and getting income. To understand more the some soil chemical properties of tidal swamp after 30 years ago opening of these lands, it will be discussed in the next section.

Soil pH, H⁺ and Al³⁺ Exchangeability

Soil acidity shown by low pH is one of the most factors affecting crop production, because some nutrients are difficult to be available for plant growth in acidic condition. Low pH in acid sulfate soil is due to natural process and/or due to roots that produce organic acids leading to low pH value (Tuyent et al., 2006). Acidic condition enhances the presence of trivalence cations (Al³⁺), which are the most toxic of Al to plant (Kochian et al., 2005). In this tidal swamp showed that the soil pH level is below 5,0 after 30 years old cultivation with a value of pH ranging from 4.0-4.59 (very acidic up to acidic) in all different land typologies in combination with high exchangeable H⁺ and Al³⁺ (Table 2). The value of H⁺ and Al³⁺ exchangeability were ranging from 0.65-1.0 Cmol(+) kg⁻¹ and 1.77-4.49 Cmol(+)kg⁻¹ respectively.

Lowest pH in combination with highest H⁺ and Al³⁺ exchangeability were found in A land typology after rice growing (Table 2), it could be due to nitrification process from nitrogen fertilization causing decrease soil pH. The comparable result was found in acid sulfate soil derived from Merbok, Kedah, Peninsular, Malaysia that the soil had 4.26 Cmol(+)kg⁻¹ Al³⁺ exchangeability (Elisa et al., 2016). In this regards, this low pH should be corrected it in order to become neutral. Sagala (2010) reported that soil pH derived from Banyu Urip had a value of 4.6, and after applying lime

with adosage of 2.5 ton ha⁻¹ increased soil pH up to 5.36. Other experiment showed that longterm NPK fertilization decreased soil pH about 0.54 unit (Czarnecki and Düring, 2015).

Organic Carbon (C_{org}) is an important chemical constituent of organic matter in a soil. Organic matter has the ability to reduce Al toxicity in the soil via chelation mechanism (Shamsuddin et al., 1994). High C_{org} and CEC were found in four different typologies of swamp land both in Mulia Sari and Banyu Urip (Table 3). The range of C_{org} was obtained about 44.11-62.57 g kg⁻¹, meanwhile the CEC was found with a value of 23.20-30.45 Cmol(+)kg⁻¹. Similar result was found by Sagala (2010) and Sagala et al. (2011) that acid sulfate soil from Banyu Urip was having 62.0 g kg⁻¹ C_{org}.

The C_{org} is the most important factor for soil CEC, because soil component known to contribute to soil CEC are organic matter and clay and showing linear regression. The linear regression model based on soil organic carbon was used to predict soil CEC (Rashidi and Seilsepour, 2008). Based on these variables showed that this acidic soil has potential good fertility but low pH, meanwhile the low pyrite content in the arable layer of four different typologies were obtained between 0.22-0.39% (Table 3). Shamsudin et al. (1994) found that C_{org} of acid sulfate soils derived from Malaysia were ranging from 41.9-74.9 g kg⁻¹ in a combination of pyrite between 0.02-0.60%. Czarnecki and Düring (2015) reported that NPK fertilization in long-term treatment had given significant different on C_{org} and CEC compared to control and these variables were increased. The high CEC of these soil shows that the soil has great agricultural potential if it was not for the presence of pyrite. Keeping the soil under anaerobic condition all the

Table 2: Soil pH, H⁺ and Al³⁺ Exchangeability

No	Land Typologies	Soil pH (1:1)	H ⁺	Al ³⁺
			Exchangeability (Cmol(+) kg ⁻¹)	
1	A	4.05 + 0.07 (va)	1.00 + 0.16	4.49 + 0.66
2	B	4.59 + 0.22 (a)	0.65 + 0.13	2.27 + 0.13
3	C	4.44 ± 0.09 (va)	0.75 ± 0.19	2.03 ± 0.55
4	D	4.55 ± 0.08 (a)	0.87 ± 0.10	1.77 ± 0.36

Note: va: very acidic, a: acidic.

Table 3: Organic Carbon, Cation Exchange Capacity and Pyrite Content

No	Land Typologies	Organic Carbon (g/kg)	Cation Exchange Capacity (Cmol (+)/kg)	Pyrite (%)
1	A	51.97+9.77 (vh)	30.45 + 7.53 (h)	0.39±0.18
2	B	62.56 ± 6.52 (vh)	24.65 ± 6.64 (h)	0.22+0.03
3	C	44.11 + 7.11 (h)	23.20 + 2.511 (m)	0.31±0.05
4	D	55.43 ± 6.08 (vh)	30.45 ± 0.00 (h)	0.25±0.03

Note: vh: very high, h: high, m: moderate.

time is probably a logical practice under the present circumstances, even though the choice of crops is limited (Shamsuddin and Senik, 1978).

The fluctuation of basic calcium such as Na⁺, Ca²⁺ and Mg²⁺exchangeabilities were presented in Table 4. Very low amount of Ca²⁺ was obtained in all typologies with values of 0.79-1.83 Cmol(+)⁻¹kg⁻¹. Lowest amount of Ca²⁺ and Mg²⁺exchangeability were found in A land typology, it is due to possibility leaching out of these metals from the arable layer by flooding. Meanwhile, Mg²⁺was found ranging from very low to low content with avalue of 0.15-0.90 Cmol(+)⁻¹kg⁻¹ (Table 4). Tuyen *et al.* (2006) reported that application of NPK fertilizer on long-term of paddy soil about 17 years old did not give any significant effect on Na⁺, Ca²⁺ and Mg²⁺ exchangeability compared to no fertilizer (control). Very high amount of exchangeable cation with values of 0.80-1.99 Cmol(+)⁻¹kg⁻¹ was only found on Na⁺ exchangeability in A and B typologies condition, it means that this pedogeneis of acid sulfate soil was really affected by saline water from the sea entering the tidal land during tide condition.

Very low total N was found in four different typologies with values of 1.48-1.67 gkg⁻¹ (Table

Table 4: Some Basic Cations of Natrium, Calcium and Magnesium Exchangeability

No	Land Typologies	Na ⁺	Ca ²⁺	Mg ²⁺
		Exchangeability(Cmol (+)/kg)		
1	A	0.80+ 0.27 (h)	0.79+ 0.17 (vl)	0.15+0.06 (vl)
2	B	0.69±0.16 (m)	1.83± 0.17 (vl)	0.44± 0.04 (l)
3	C	1.99+1.13 (vh)	1.77+ 0.19 ((vl)	0.90+ 0.53 (l)
4	D	0.24±0.04 (l)	1.70± 0.27 (vl)	0.54 ± 0.17 (l)

Note: vl: very low, l: low, m: moderate, h: high, vh: very high.

Table 5: Macro Nutrients of N, P and K in Tidal Swamp Land

No	Land Typologies	Total N (g/kg)	P (mg/kg)	K Cmol(+)/kg
1	A	1.68+ 0.93 (l)	9.40 +2.32 (vl)	0.28± 0.13 (l)
2	B	1.87± 0.07 (l)	71.35±23.19 (vh)	0.79± 0.48 (h)
3	C	1.48+ 0.27 (l)	70.25+ 7.88 (vh)	0.32± 0.00 (l)
4	D	1.68± 0.24 (l)	58.95±12.54 (vh)	0.58± 0.08 (h)

Note: vl: very low, l: low, h: high, vh: very high.

5) and very low of P-Bray I was only obtained in A typology, other three typologies were found P in high content. Furthermore, K exchangeability was obtained ranging from low to high with a value of 0.28-0.79 Cmol(+)⁻¹ kg (Table 5). The low total N content after along term of tidal swamp cultivation for food crop was due to low pH which did not give any microbe chance to do mineralization of organic matter.

CONCLUSION

Tidal swamp land after 30 years opened, still have high organic carbon and high cation exchange capacity but low pH, and as well as low calcium,

magnesium and total nitrogen respectively. Relatively, high content of nutrient was only found for Na and some P in soil. These tidal lands have potential to be cultivated for agriculture after improving soil pH and alleviating Al and absence of pyrite from arable layer.

ACKNOWLEDGMENT

The financial support provided by the Universitas Sriwijaya under excellent research professor profession award number 1023/UN9.3.1/LPPM/2016 is gratefully acknowledged

REFERENCES

- Bertsch P M and Bloom P R (1996), "Aluminum", in Bartels J M (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 517-550.
- Bremner J M (1996), "Nitrogen-Total", in Bartels J M (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 1085-1122.
- Budianta D and Windusari Y (2016), "Beneficial Effect of Local Resources to Improve Food Crop Production in Tidal Swamp of Indonesia", *Inter. J. Env & Agric Res*, Vol. 2, No. 1, pp. 98-101.
- Budianta D, Napoleon A and Ristiani D (2012), SRI di Lahan Pasang Surut, Prosiding Seminar Nasional: Menuju Pertanian Berdaulat, Tanggal 12 September 2012, Fakultas Pertanian Universitas Bengkulu bekerjasama dengan PERHEPI dan PFI Komda Bengkulu, Hal, pp. 19-25.
- Czarnecki S and Dûring R A (2015), "Influence of Long-Term Mineral Fertilization on Metal Contents and Properties of Soil Samples Taken from Different Locations in Hesse, Germany", *Soil*, Vol. 1, pp. 23-33.
- Dent D (1986), "Acid Sulphate Soils: Baseline for Research and Development", p. 204, *ILRI*, Wageningen. Publ. No. 39, The Netherland.
- Dublin-Green C O and Ojanuga A G (1988), "The Problem of Acid Sulfate Soils in Brcaish Water Aquaculture: A Preliminary Study of The Soils of Niomr/Atac Fish Farm, Buguma, Rivers State, Nigeria", Nigerian Institute for Oceanography and Marine Research, Technical Paper No. 45, November.
- Elisa AA, Shamsuddin J and Roslan I (2016), "Alleviating Aluminum Toxicity in an Acid Sulfate Soil from Peninsular, Malaysia by Calsium Silicate Application", *Solid Earth*, Vol. 7, pp. 367-374.
- Euroconsult (1995), Laporan Pemantauan Aspek-aspek Hidrologi Makro: Proyek Pengembangan Pertanian Telang dan Saleh, Komponen Pengembangan Drainase, Integrated Irrigation Sector Project (IISP) (In Indonesian).
- Fahmi A and Hanudin E (2008), Pengaruh Kondisi Redoksterhadap Stabilitas Kompleks Organik-besipada Tanah SulfatMasam", *J. Ilmu Tanah danLingkungan*, Vol. 8, No. 1, pp. 49-55.
- Hairani A, Yusuf A, Purnomo E and Osaki M (2005), Identifikasi dan Uji Kemampuan Mikroorganisme Pelarut P dari Berbagai Rhizosfer Padi Lokal yang Berdaya hasil Tinggi DI Lahan Pasang Surut Tanah Sulfat masam Kp. Belandean Kalimantan Selatan, *J. Tanah Trop.*, Vol. 10, No. 2, pp. 131-136.
- Helmke P A and Sparks D L (1996), "Lithium, Sodium, Potasium, Rubidium and Cesium,

- in Bartels J M (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 551-574.
13. Ilona N A, Munif G, Sopandie D, Sutandi A and Melati M (2012), "The Utilization of Acid Sulfate Soil for Agrivulture", in *International Seminar: Food Sovereignty and Natural Resources in Archipelago Region*, October 23-24, pp. 261-270, ICC-IPB.
 14. Imanudin M S, Armanto E, Susanto R H and Bernas S M (2010), "Water Table Fluctuation in Tidal Lowland for Developing Agricultural Water Management Strategies", *J. Trop. Soils*, Vol. 15, No. 3, pp. 277-282.
 15. Intimulya Multikencana (2009), "Review Desain Daerah Rawa Pasang Surut Delta Telang II", Kabupaten Banyuasin, Propinsi Sumatera Selatan, Final Report.
 16. Kochian L V, Pineros M A and Hoekenga O A (2005), "The Physiology, Genetics and Molecular Biology of Plant Aluminum Resistance and Toxicity", *Plant Soil*, Vol. 274, pp. 175-195.
 17. Koesrini E William and Nursyamsi D (2014), "Application of Lime and Adaptable Variety to Increase Tomato Productivity at Potential Acid Sulphate Soil", *J. Trop Soils*, Vol. 19, No. 2, pp. 69-76.
 18. Koesrini M Saleh and Nursyamsi D (2013), Keragaan Varietas Inpara di Lahan Rawa Pasang Surut (*Performance of Varieties Inpara in Swamp Land*), *Pangan*, Vol. 22, No. 3, pp. 221-228.
 19. Koswara O and Rumawas F (1984), Tidal Swamp Rice in Palembang Region, in *Proceedings Workshop on Research Priorities in Tidal Samp Rice*, pp. 37-46, IRRl, Los Banos, Philippines.
 20. Kuo S (1996), "Phosphorus", in Bartels J M (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 869-920.
 21. Marlina L, Imanudin M I and Priatna S J (2015), *Kajian Pola Pemanfaatan Lahan di Daerah Reklamasi Pasang Surut Delta Telang II Kabupaten Banyuasin (The Study of Land Use Pattern in Tidal Lowland Reclamations of Telang II Deltaic Areas of Banyuasin Reegency)*, in *Prosiding Seminar Nasional Dalam Rangka Dies Natalis k3-52 Fakultas Pertanian Universitas Sriwijaya*, Palembang, November 5.
 22. Masulili A, Utomo W H and Wisnubroto E I (2016), "Growing Rice (*Orizasetiva*L) in the Sulphate Acid Soils of West Kaliman, Indonesia", *Inter. J. Agric. Res*, Vol. 11, No. 1, pp. 13-22.
 23. Moore Jr P A, Attanandana T and Pattrick Jr W H (1990), "Factors Affeting Rice Growth on Acid Sulfate Soils", *Soil Sci. Soc. Am. J*, Vol. 54, pp. 1651-1656.
 24. Muktamar Z and Adiprasetyo T (1993), *Studi Potensi Lahan Gambut di Propinsi Bengkulu Untuk Tanaman Semusim*, *Prosiding Seminar Nasional Gambut II*, Bengkulu.
 25. Nelson D W and Sommers L E (1996), "Total Carbon, Organic Carbon and Organic Matter", in Bartels J M (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 961-1010.
 26. Nursanti I (2014), *Peranan Mineral Zeolit dan produk Samping Cair Pabrik Kelapa Sawit terhadap Pertumbuhan dan Produksi Tanaman Jagung pada Keadaan Cekaman Kekeringan di Tanah Sulfat Masam Potensial*, Disertasi FP, Unsri, 161 hal.

27. Purnomo E, Hasegawa T, Hashidoko Y and Osaki M (2006), "Soil Nitrogen Supply and Nitrogen Uptake for Local Rice Grown in Unfertilized Acid Sulfate Soil in South Kalimantan", *Tropics*, Vol. 15, No. 4, pp. 349-354.
28. Rahmi O, Susanto R H and Siswanto A (2015), Pengelolaan Lahan Basah Terpadu di Desa Mulia Sari Kecamatan Tanjung Lago, Kabupaten Banyuasin (*The Integrated Lowland Management in Mulia Sari, Tanjung Lago Sybdistrict, Banyuasin Regency*), *J. I. Pert. Indo (JIPI)*, Vol. 20, No. 3, pp. 201-207.
29. Rashidi M and Seilsepour M (2008), "Modeling of Soil Cation Exchange Capacity Based on Soil Organic Carbon", *ARPN J. Agric. & Bio. Sci.*, Vol. 3, No. 4, pp. 41-45.
30. Ristiani D (2012), "Modifikasi System of Rice Intensification (SRI) di Lahan Pasang Surut Dengan Berbagai Umur Pindah Bibit dan Pemberian Pupuk Organik Cair, Thesis S2, Universitas Sriwijaya.
31. Sagala D (2010), Peningkatan pH Tanah Masam di Lahan Rawa Pasang Surut pada Berbagai Dosis Kapur Untuk Budidaya Kedelai (*The Increase of pH of Acid Tidal Swamps on Dosage of Liming for Soybean Culture*), *J. Agroqua*, Vol. 8, No. 2, pp. 1-5.
32. Sagala D, Ghulamahdi dan M and Melati M (2011), Pola Serapan Hara dan Pertumbuhan Beberapa Varietas Kedelai dengan Budidaya Jenuh Air di Lahan Rasa Pasang Surut, *J. Agroqua*, Vol. 9, No. 1, pp. 1-10.
33. Shamsuddin J and Senik T O R (1978), "Mineralogy and Related Properties of Acid Sulphate Soil from Melaka, Peninsular Malaysia", *Pertanika*, Vol. 1, No. 2, pp. 136-142.
34. Shamsuddin J, Jamilah I and Ogunwale JA (1994), "Organic Carbon Determination in Acid Sulphate Soils", *Pertanika J. Trop. Agric. Sci*, Vol. 17, No. 3, pp. 197-200.
35. Suarez D L (1996), "Beryllium, Magnesium, Calcium, Strontium and Barium", in Bartels JM (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 575-602.
36. Sumner M E and Miller W/P (1996), "Cation Exchange Capacity and Exchange Coefficients", in Bartels JM (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 1201-1230.
37. Tabatabai MA (1996), "Sulfur", in Bartels JM (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 921-960.
38. Thomas G W (1996), "Soil pH and Soil Acidity", in Bartels J M (1996), "Methods of Soil Analysis: Part 3 Chemical Methods", *Soil Sci. Ame. Book Series*, pp. 475-490.
39. Tuyent T Q, Phung C V and Tiuh T K (2006), "Influence of Long Term Application of N, P, K Fertilization on Soil pH, Organic Matter, CEC, Exchangeable Cations and Some Trace Elements", *Omonrice*, Vol. 14, pp. 144-148.
40. Widjaja Adhi I P G, Nugroho K, Ardi A and Karama A S (1992), Sumberdaya Lahan Rawa: Potensi, Keterbatasan dan Pemanfaatan", Prosiding: Pengembangan Terpadu Pertanian Lahan Rawa Pasang Surut dan Lebak.

41. Wijanarko A, Taufik A and Harnowo D (2016), "Effect of Liming, Manura, and NPK Fertilizer Application on Growth and Yield Performance of Soybean in Swam Land", *J. Degraded and Mining Lands Manag*, Vol. 3, No. 2, pp. 527-533.



International Journal of Engineering Research and Science & Technology

Hyderabad, INDIA. Ph: +91-09441351700, 09059645577

E-mail: editorijerst@gmail.com or editor@ijerst.com

Website: www.ijerst.com

